

POSTER PRESENTATION

Open Access

Tumor-infiltrating T lymphocyte clonality predicts prognosis in human ovarian cancer

Takemasa Tsuji^{1*}, Kevin H Eng², Sacha Gnjatich³, Junko Matsuzaki¹, Rachel Brightwell⁴, Anthony Miliotto¹, Ryan Emerson⁵, Cindy Desmarais⁵, Erika Lindsley⁵, Julie Rubinstein⁵, Jianmin Wang², Song Liu², Harlan Robins⁵, Kunle Odunsi^{1,4}

From Society for Immunotherapy of Cancer 28th Annual Meeting
National Harbor, MD, USA. 8-10 November 2013

Background

The prognostic significance of the number of tumor-infiltrating T cells has been demonstrated for many tumor types. In contrast, the significance of the tumor-infiltrating T cell clonality, which reflects the preferential infiltration or expansion of T cell clones in the tumor microenvironment, has not been clear because of the technical hurdles required for evaluating each T cell clone in the tumor. In order to delineate the complexity of T cell responses and define correlates of a protective immunity, we applied a recently developed deep T cell receptor (TCR)-sequencing technology (immunoSEQ) to paired frozen tumor tissues and peripheral blood mononuclear cells from 99 ovarian cancer patients.

Methods

Rearranged β TCR chain DNA sequences were sequenced using immunoSEQ technology. Frequency of each T cell clone was obtained from the copy number of the sequence. T cell clonality of the specimens was calculated from entropy of TCR sequences. Spontaneous immune responses against tumor-associated antigens (NY-ESO-1, MAGE-A1, MAGE-A3 and p53) were evaluated by measuring serum antibodies by ELISA.

Results

Approximately $3(\pm 3) \times 10^6$ and $4(\pm 2) \times 10^6$ full-length TCR beta chain sequences were obtained corresponding to the detection limit for T cell frequency at 3×10^{-7} and 2×10^{-7} for tumor and blood samples, respectively. In patients who had spontaneous antibodies against a panel of tumor-associated antigens, more clonal T cell infiltration

was associated with longer progression-free survival. In sharp contrast, clonal infiltration was a worse prognostic factor in patients without detectable humoral immune responses against surrogate tumor antigens. From sequence based analyses, we found a set of shared TCR sequences among patients.

Conclusion

Deep TCR sequencing using immunoSEQ technology is a powerful tool to characterize tumor-infiltrating T cell clonality using frozen tumor tissues. Our analyses indicate that evaluation for spontaneous anti-tumor immune responses is required to correctly understand the prognostic significance of tumor-infiltrating T cells.

Authors' details

¹Center for Immunotherapy, Roswell Park Cancer Institute, Buffalo, NY, USA. ²Biostatistics and Bioinformatics, Roswell Park Cancer Institute, Buffalo, NY, USA. ³Tisch Cancer Institute, Icahn School of Medicine at Mount Sinai, New York, NY, USA. ⁴Gynecologic Oncology, Roswell Park Cancer Institute, Buffalo, NY, USA. ⁵Adaptive Biotechnologies, Seattle, WA, USA.

Published: 7 November 2013

doi:10.1186/2051-1426-1-S1-P64

Cite this article as: Tsuji et al.: Tumor-infiltrating T lymphocyte clonality predicts prognosis in human ovarian cancer. *Journal for ImmunoTherapy of Cancer* 2013 **1**(Suppl 1):P64.

¹Center for Immunotherapy, Roswell Park Cancer Institute, Buffalo, NY, USA
Full list of author information is available at the end of the article